

ON THE CALCULATION OF GROUP CHARACTERS

M. Gungormez

Dept. Physics, Fac. Science, Istanbul Tech. Univ.
80626, Maslak, Istanbul, Turkey
e-mail: gungorm@itu.edu.tr

H. R. Karadayi

Dept. Physics, Fac. Science, Istanbul Tech. Univ.
80626, Maslak, Istanbul, Turkey
e-mail: karadayi@itu.edu.tr

and

Dept. Physics, Fac. Science, Istanbul Kultur University
34156, Atakoy, Istanbul, Turkey

Abstract

It is known that characters of irreducible representations of finite Lie algebras can be obtained by Weyl character formula including Weyl group summations which make actual calculations almost impossible except for a few Lie algebras of lower rank. By starting from Weyl character formula, we show that these characters can be re-expressed without referring to Weyl group summations. Some useful technical points are given in detail in the instructive example of G_2 Lie algebra.

I. INTRODUCTION

It is known that the character formula of Weyl [1] gives us a direct way to calculate the character of irreducible representations of finite Lie algebras. For this, let G_r be a Lie algebra of rank r , $W(G_r)$ its Weyl group, α_i 's and λ_i 's be, respectively, its simple roots and fundamental dominant weights. The notation here and in the following sections will be as in our previous work [2]. For further reading, we refer to the beautiful book of Humphreys [3]. A dominant weight Λ^+ is expressed in the form

$$\Lambda^+ = \sum_{i=1}^r s_i \lambda_i \quad (I.1)$$

where s_i 's are some positive integers (including zero). An irreducible representation $V(\Lambda^+)$ can then be attributed to Λ^+ . The character $Ch(\Lambda^+)$ of $V(\Lambda^+)$ is defined by

$$Ch(\Lambda^+) \equiv \sum_{\lambda^+} \sum_{\mu \in W(\lambda^+)} m_{\Lambda^+}(\mu) e^\mu \quad (I.2)$$

where $m_{\Lambda^+}(\mu)$'s are multiplicities which count the number of times a weight μ is repeated for $V(\Lambda^+)$. The first sum here is over Λ^+ and all of its sub-dominant weights λ^+ 's while the second sum is over the elements of their Weyl orbits $W(\lambda^+)$'s. Formal exponentials are taken just as in the book of Kac [4] and in (I.2) we extend the concept for any weight μ , in the form e^μ . Note here that multiplicities are invariant under Weyl group actions and hence it is sufficient to determine only $m_{\Lambda^+}(\lambda^+)$ for the whole Weyl orbit $W(\lambda^+)$.

An equivalent form of (I.2) can be given by

$$Ch(\Lambda^+) = \frac{A(\rho + \Lambda^+)}{A(\rho)} \quad (I.3)$$

where ρ is the Weyl vector of G_r . (I.3) is the celebrated **Weyl Character Formula** which gives us the possibility to calculate characters in the most direct and efficient way. The central objects here are $A(\rho + \Lambda^+)$'s which include a sum over the whole Weyl group:

$$A(\rho + \Lambda^+) \equiv \sum_{\sigma \in W(G_r)} \epsilon(\sigma) e^{\sigma(\rho + \Lambda^+)} . \quad (I.4)$$

In (I.4), σ denotes an element of Weyl group, i.e. a Weyl reflection, and $\epsilon(\sigma)$ is the corresponding **signature** with values either +1 or -1.

The structure of Weyl groups is completely known for finite Lie algebras in principle. In practice, however, the problem is not so trivial, especially for Lie algebras of some higher rank. The order of E_8 Weyl group is, for instance, 696729600 and any application of Weyl character formula needs for E_8 an explicit calculation of a sum over 696729600 Weyl reflections. Our main point here is to overcome this difficulty in an essential manner.

For an actual application of (I.3), an important notice is the **specialization** of formal exponentials e^μ as is called in the book of Kac [4]. In its most general form, we consider here the specialization

$$e^{\alpha_i} \equiv u_i, \quad i = 1, 2, \dots, r. \quad (I.5)$$

which allows us to obtain $A(\rho)$ in the form of

$$A(\rho) = P(u_1, u_2, \dots, u_r) \quad (I.6)$$

where $P(u_1, u_2, \dots, u_r)$ is a polynomial in indeterminates u_i 's. We also have

$$A(\rho + \Lambda^+) = P(u_1, u_2, \dots, u_r; s_1, s_2, \dots, s_r) \quad (I.7)$$

where $P(u_1, u_2, \dots, u_r; s_1, s_2, \dots, s_r)$ is another polynomial of indeterminates u_i 's and also parameters s_i 's defined in (I.1). The Weyl formula (I.3) then says that polynomial (I.7) always factorizes on polynomial (I.6) leaving us with another polynomial $R(u_1, u_2, \dots, u_r; s_1, s_2, \dots, s_r)$ which is nothing but the character polynomial of $V(\Lambda^+)$.

The specialization (I.5) will always be normalized in such a way that (I.3) gives us the **Weyl dimension formula** [5], in the limit $u_i = 1$ for all $i = 1, 2, \dots, r$. One also expects that

$$P(u_1, u_2, \dots, u_r; 0, 0, \dots, 0) \equiv P(u_1, u_2, \dots, u_r). \quad (I.8)$$

II. RECREATING $A(\rho + \Lambda^+)$ FROM $A(\rho)$

In this section, without any reference to Weyl groups, we give a way to calculate polynomial (I.7) directly from polynomial (I.6). For this, we first give the following explicit expression for polynomial (I.6):

$$A(\rho) = \frac{\prod_{\alpha \in \Phi^+} (e^\alpha - 1)}{\prod_{i=1}^r (e^{\alpha_i})^{k_i}} \quad (II.1)$$

where

$$k_i \equiv \frac{1}{2} (\alpha_i, \alpha_i) (\lambda_i, \rho) \quad (II.2)$$

and Φ^+ is the positive root system of G_r . Exponents k_i 's are due to the fact that the monomial of maximal order is

$$\prod_{i=1}^r (e^{\alpha_i})^{2k_i}$$

in the product $\prod_{\alpha \in \Phi^+} (e^\alpha - 1)$. All the scalar products like (λ_i, α_j) or (α_i, α_i) are the symmetrical ones and they are known to be defined via Cartan matrix of a Lie algebra. The crucial point, however, is to see that (II.1) is equivalent to

$$A(\rho) = \prod_{A=1}^{|W(G_r)|} \epsilon_A (e^{\alpha_i})^{\xi_i^0(A)} \quad (II.3)$$

where $|W(G_r)|$ is the order of Weyl group $W(G_r)$ and as is emphasized in section I, ϵ_A 's are signatures with values $\epsilon_A = \mp 1$. Note here that, by expanding the product $\prod_{\alpha \in \Phi^+} (e^\alpha - 1)$, the precise values of signatures can be determined uniquely.

To get an explicit expression for exponents $\xi_i^0(A)$ in (II.3), let us define R^+ is composed out of elements of the form

$$\beta^+ \equiv \sum_{i=1}^r n_i \alpha_i \quad (II.4)$$

where n_i 's are some positive integers including zero. R^+ is a subset of the Positive Root Lattice of G_r . The main emphasis here is on some special roots

$$\gamma_i(I_i) \in R^+$$

which are defined by following conditions

$$(\lambda_i - \gamma_i(I_i), \lambda_j - \gamma_j(I_j)) = (\lambda_i, \lambda_j) \quad , \quad i, j = 1, 2, \dots, r. \quad (II.5)$$

Note here that (λ_i, λ_j) 's are defined by inverse Cartan matrix. For the range of indices I_j 's, we assume that they take values from the set $\{1, 2, \dots, |I_j|\}$, that is

$$I_j \in \{1, 2, \dots, |I_j|\} \quad , \quad j = 1, 2, \dots, r.$$

We also define the sets

$$\Gamma(A) \equiv \{\gamma_1(I_1(A)), \gamma_2(I_2(A)), \dots, \gamma_r(I_r(A))\} \quad , \quad A = 1, 2, \dots, D \quad (II.6)$$

are chosen by conditions (II.5) where $I_j(A) \in \{1, 2, \dots, |I_j|\}$ and D is the maximal number of these sets.

Following two statements are then valid:

$$(1) \quad \mathbf{D} = |\mathbf{W}(\mathbf{G}_r)|$$

$$(2) \quad |\mathbf{I}_j| = |\mathbf{W}(\lambda_j)|$$

where $|W(\lambda_j)|$ is the order, i.e. the number of elements, of the Weyl orbit $W(\lambda_j)$. As is known, a Weyl orbit is stable under Weyl reflections and hence all its elements have the same length. It is interesting to note however that the lengths of any two elements $\gamma_i(j_1)$ and $\gamma_i(j_2)$ could, in general, be different while the statement (2) is, still, valid.

The exponents in (II.3) can now be defined by

$$\xi_i^0(A) \equiv \frac{1}{2} (\alpha_i, \alpha_i) (\lambda_i - \gamma_i(A), \rho) \quad . \quad (II.7)$$

We then also state that the natural extension of (II.3) is as in the following:

$$A(\rho + \Lambda^+) = \prod_{A=1}^{|W(G_r)|} \epsilon_A (e^{\alpha_i})^{\xi_i(A)} \quad (II.8)$$

for which the exponents are

$$\xi_i(A) \equiv \frac{1}{2} (\alpha_i, \alpha_i) (\lambda_i - \gamma_i(A), \rho + \Lambda^+) \quad . \quad (II.9)$$

as a natural extension of (II.7). Signatures have the same values in both expressions, (II.3) and (II.8). This reduces the problem of explicit calculation of character polynomial $Ch(\Lambda^+)$ to the problem of finding solutions to conditions (II.5). It is clear that this is more manageable than that of using Weyl character formula directly.

III. AN EXAMPLE : $\mathbf{G}_2$

It is known that G_2 is characterized by two different root lengths

$$(\alpha_1, \alpha_1) = 6 \quad , \quad (\alpha_2, \alpha_2) = 2 \quad (III.1)$$

and also

$$(\alpha_1, \alpha_2) = -3 \quad . \quad (III.2)$$

Its positive root system is thus

$$\Phi^+ = \{ \alpha_1 , \alpha_2 , \alpha_1 + \alpha_2 , \alpha_1 + 2\alpha_2 , \alpha_1 + 3\alpha_2 , 2\alpha_1 + 3\alpha_2 \} \quad (III.3)$$

Beyond Φ^+ , first few elements of R^+ are determined by

$$\begin{aligned} & 2\alpha_2 , \quad 2\alpha_1 + 2\alpha_2 , \quad 2\alpha_1 + 4\alpha_2 , \quad \alpha_1 + 4\alpha_2 , \\ & 2\alpha_1 + \alpha_2 , \quad 2\alpha_1 + 5\alpha_2 , \quad 3\alpha_1 + 4\alpha_2 , \quad 3\alpha_1 + 5\alpha_2 , \quad 3\alpha_2 , \\ & 3\alpha_1 + 3\alpha_2 , \quad 3\alpha_1 + 6\alpha_2 , \quad 2\alpha_1 , \quad 2\alpha_1 + 6\alpha_2 , \quad 4\alpha_1 + 6\alpha_2 \end{aligned}$$

as will be seen from the definition (II.4). This will be sufficient to obtain following elements which fulfill conditions (II.5):

$$\begin{aligned} \gamma_1(1) &= 0 , \\ \gamma_1(2) &= \alpha_1 , \\ \gamma_1(3) &= \alpha_1 + 3\alpha_2 , \\ \gamma_1(4) &= 3\alpha_1 + 3\alpha_2 , \\ \gamma_1(5) &= 3\alpha_1 + 6\alpha_2 , \\ \gamma_1(6) &= 4\alpha_1 + 6\alpha_2 \end{aligned}$$

and

$$\begin{aligned} \gamma_2(1) &= 0 , \\ \gamma_2(2) &= \alpha_2 , \\ \gamma_2(3) &= \alpha_1 + \alpha_2 , \\ \gamma_2(4) &= \alpha_1 + 3\alpha_2 , \\ \gamma_2(5) &= 2\alpha_1 + 3\alpha_2 , \\ \gamma_2(6) &= 2\alpha_1 + 4\alpha_2 \end{aligned}$$

from which we obtain, as in (II.6), following 12 sets of 2-elements:

$$\begin{aligned}
\Gamma_1 &= \{\gamma_1(1), \gamma_2(1)\} \quad , \quad \epsilon_1 = +1 \\
\Gamma_2 &= \{\gamma_1(2), \gamma_2(3)\} \quad , \quad \epsilon_2 = +1 \\
\Gamma_3 &= \{\gamma_1(3), \gamma_2(2)\} \quad , \quad \epsilon_3 = +1 \\
\Gamma_4 &= \{\gamma_1(4), \gamma_2(5)\} \quad , \quad \epsilon_4 = +1 \\
\Gamma_5 &= \{\gamma_1(5), \gamma_2(4)\} \quad , \quad \epsilon_5 = +1 \\
\Gamma_6 &= \{\gamma_1(6), \gamma_2(6)\} \quad , \quad \epsilon_6 = +1 \\
\Gamma_7 &= \{\gamma_1(1), \gamma_2(2)\} \quad , \quad \epsilon_7 = -1 \\
\Gamma_8 &= \{\gamma_1(2), \gamma_2(1)\} \quad , \quad \epsilon_8 = -1 \\
\Gamma_9 &= \{\gamma_1(3), \gamma_2(4)\} \quad , \quad \epsilon_9 = -1 \\
\Gamma_{10} &= \{\gamma_1(4), \gamma_2(3)\} \quad , \quad \epsilon_{10} = -1 \\
\Gamma_{11} &= \{\gamma_1(5), \gamma_2(6)\} \quad , \quad \epsilon_{11} = -1 \\
\Gamma_{12} &= \{\gamma_1(6), \gamma_2(5)\} \quad , \quad \epsilon_{12} = -1
\end{aligned} \tag{III.4}$$

Note here that $|W(G_2)| = 12$, $|W(\lambda_1)| = 6$ and $|W(\lambda_2)| = 6$ show, for G_2 , the validity of our two statements mentioned above. The signatures are given on the right-hand side of (III.4). They can be easily found to be true by comparing two expressions (II.1) and (II.3) for the case in hand. As in (I.5), let us choose the most general specialization

$$e^{\alpha_1} = x \quad , \quad e^{\alpha_2} = y \quad .$$

Then, in view of (I.8), character polynomial

$$R(x, y, s_1, s_2) \equiv \frac{P(x, y, s_1, s_2)}{P(x, y, 0, 0)} \tag{III.5}$$

for an irreducible G_2 representation originated from a dominant weight $\Lambda^+ = s_1\lambda_1 + s_2\lambda_2$ will be expressed in its following **ultimate** form:

$$\begin{aligned}
P(x, y, s_1, s_2) &= + x^{3+2 \ s_1+s_2} y^{5+3 \ s_1+2 \ s_2} + x^{-3-2 \ s_1-s_2} y^{-5-3 \ s_1-2 \ s_2} \\
&+ x^{-2-s_1-s_2} y^{-1-s_2} + x^{2+s_1+s_2} y^{1+s_2} \\
&+ x^{-1-s_1} y^{-4-3 \ s_1-s_2} + x^{1+s_1} y^{4+3 \ s_1+s_2} \\
&- x^{1+s_1} y^{-1-s_2} - x^{-1-s_1} y^{1+s_2} \\
&- x^{-2-s_1-s_2} y^{-5-3 \ s_1-2 \ s_2} - x^{2+s_1+s_2} y^{5+3 \ s_1+2 \ s_2} \\
&- x^{-3-2 \ s_1-s_2} y^{-4-3 \ s_1-s_2} - x^{3+2 \ s_1+s_2} y^{4+3 \ s_1+s_2}
\end{aligned} \tag{III.6}$$

In the following we will give some examples to illustrate (III.5);

$$Ch(\lambda_1) = R(x, y, 1, 0) = \frac{1}{x^2 y^3} (1 + x + x y + x y^2 + x^2 y^2 + x y^3 + 2 x^2 y^3 + x^3 y^3 \\ + x^2 y^4 + x^3 y^4 + x^3 y^5 + x^3 y^6 + x^4 y^6)$$

$$Ch(\lambda_2) = R(x, y, 0, 1) = \frac{1}{x y^2} (1 + y + x y + x y^2 + x y^3 + x^2 y^3 + x^2 y^4)$$

$$Ch(\lambda_1 + \lambda_2) = R(x, y, 1, 1) = \frac{1}{x^3 y^5} (1 + x) (1 + y) (1 + x y) (1 + x y^2) (1 + \\ x y^3) (1 + x^2 y^3)$$

$$Ch(2 \lambda_2) = R(x, y, 0, 2) = \frac{1}{x^2 y^4} (1 + y + y^2) (1 + x y + x^2 y^2) (1 + x y^2 + x^2 y^4)$$

where

$$P(x, y, 0, 0) = \frac{1}{x^3 y^5} (1 - x) (1 - y) (1 - x y) (1 - x y^2) (1 - x y^3) (1 - x^2 y^3)$$

A useful application of (III.5) is in the calculation of tensor coupling coefficients [6]. We note, for instance, that the coefficients in decomposition

$$V(\lambda_1) \otimes V(\lambda_1 + \lambda_2) = V(2 \lambda_1 + \lambda_2) \oplus V(\lambda_1 + 2 \lambda_2) \oplus V(4 \lambda_2) \oplus V(3 \lambda_2) \\ \oplus 2 V(\lambda_1 + \lambda_2) \oplus V(2 \lambda_2) \oplus V(\lambda_2)$$

can be calculated easily by

$$R(x, y, 1, 0) \otimes R(x, y, 1, 1) = R(x, y, 2, 1) \oplus R(x, y, 1, 2) \oplus R(x, y, 0, 4) \oplus R(x, y, 0, 3) \\ \oplus 2 R(x, y, 1, 1) \oplus R(x, y, 0, 2) \oplus R(x, y, 0, 1)$$

where the terms above are already known by (III.5).

CONCLUSION

Although application of our method is given here in case of G_2 Lie algebra, there are no serious difficulty to do the same for classical chains A_r , B_r , C_r and D_r of finite Lie algebras. Our results for exceptional Lie algebras F_4 and E_6 will be given soon in a subsequent publication. In case of E_7 and E_8 Lie algebras, one must however emphasize that rather than the most general specialization (I.5) of formal exponentials we need in practice more simple specializations though there are some other methods [7] which work equally well for E_7 and E_8 . For Affine Lie Algebras, a similar study of Weyl-Kac character formula will also be interesting.

REFERENCES

- [1] H.Weyl, The Classical Groups, N.J. Princeton Univ. Press (1946)
- [2] H.R.Karadayi and M.Gungormez, J.Phys.A:Math.Gen. 32 (1999) 1701-1707
- [3] J.E.Humphreys, Introduction to Lie Algebras and Representation Theory, N.Y., Springer-Verlag (1972)
- [4] V.G.Kac, Infinite Dimensional Lie Algebras, N.Y., Cambridge Univ. Press (1990)
- [5] Weyl dimension formula cited in p.139 of ref [3]
- [6] Steinberg formula cited in p.140 of ref [3]
- [7] H.R.Karadayi and M.Gungormez, Summing over the Weyl Groups of E_7 and E_8 , math-ph/9812014